

CHEMICAL EVOLUTION OF THE GALAXY: THE G-DWARF PROBLEM AND RADIOACTIVE CHRONOLOGY REVISITED TAKING ACCOUNT OF THE THICK DISK

B.E.J. PAGEL

Astronomy Centre, CPES, University of Sussex, Brighton BN1 9QJ, UK
E-mail: bejp@star.cpes.susx.ac.uk

Thin and Thick disk metallicities overlap, but $\alpha(\text{Fe})$ relations differ; there is also a hiatus in time before the first thin-disk stars, with a minimum metallicity $[\text{Fe}/\text{H}] \simeq -0.6$. Thus there are two ‘G-dwarf’ problems. We fit metallicity distribution functions (MDFs) for the two disks derived by Wyse & Gilmore (1995) with analytical models (for O and Mg). The Thick disk fits a simple inflow model with effective yield and final $[\text{Mg}/\text{H}]$ about the same as for the Thin disk, excluding continuous mass loss from the former as a source of ‘prompt initial enrichment’ (PIE) for the latter. The Thin disk has a narrow MDF fitted (rather poorly) with a closed model having PIE from some other source, or more elegantly with a slow infall model similar to the ‘two-inflow’ scenario of Matteucci et al., but with a PIE and a severe break in the age-metallicity relation. Implications for radio-active cosmochemistry are investigated.

1 Introduction

The ‘G-dwarf’ problem has been with us for nearly 40 years and extended to other galaxies. There is still no definitive universally accepted solution; rather too many, the most popular being ‘prompt initial enrichment’ (PIE; Truran & Cameron 1971) and inflow (Larson 1972). Truran & Cameron postulated a first generation of massive stars, sometimes called Population III, collapsing to black holes or ‘collapsars’ in the halo, thereby accounting for its dark mass. Later the idea gained ground that the halo consists mostly of dark matter, wholly or partly non-baryonic, but PIE came in by another route. After discovery of the Thick Disk, Gilmore & Wyse (1986) suggested that (a) the Thick disk was initially enriched by gas shed from the halo as in the model by Hartwick (1976); and (b) the Thin disk was similarly enriched by gas shed from the Thick one. I never took much notice of this paper, because the relative numbers are not right unless there is inflow as well, and Wyse & Gilmore (1992) themselves abandoned proposition (a) on angular momentum grounds. This still leaves open a connection between the two disks, and Wyse & Gilmore (1995) have published separate MDFs for them which, with new data on O/Fe and Mg/Fe, have implications that I discuss in this talk.




Figure 1. Iron MDFs for Thick and Thin disks, after Gilmore & Wyse (1995).

2 Iron and magnesium MDFs and AMRs

Numerous models based on inflow give a fair fit to the MDF of nearby disk stars as a whole (Matteucci & François 1989; Pagel 1989; Pagel & Tautvaišienė 1995). PIE models based on outflow of metals from a rapidly evolving bulge have also been quite successful (Köppen & Arimoto 1990; Samland et al. 1997). Strobel (1991), Pardi et al. (1995), Chiappini et al. (1997), Fuhrmann (1998) and Gratton et al. (2000), among others, have investigated chemical evolution implications of the different time-scales of evolution of the two disks, and Matteucci and co-authors have developed ‘two-inflow’ models which I consider a good idea, but I shall argue that they do not go far enough.




Figure 2. $[Mg/Fe]$ vs $[Fe/H]$ and $[Fe/Mg]$ vs $[Mg/H]$ for stars of the Galactic halo, thick disk and thin disk, after Fuhrmann (1998). Courtesy Klaus Fuhrmann.

Wyse & Gilmore’s iron MDFs are shown in Figure 1, where it is seen that the Thick disk has a mean $[Fe/H]$ just at the sharp lower limit of the Thin one,

which made it seem natural for Gilmore & Wyse (1986) to suggest a Hartwick-like model for the Thick disk where gas expelled from it over time led to a low effective yield and had just the right metallicity to start off the Thin disk, the mass of which can be made up as desired by subsequent inflow. This, however, assumes a single $\alpha(\text{Fe})$ relation, which now seems incorrect (Fuhrmann 1998; Gratton et al. 2000), as is illustrated in Figure 2. The Thick-disk stars have enhanced α/Fe (and O/Fe) even when their α -metallicity is as high as solar.




Figure 3. Thin curves: age-metallicity relation from the ‘two-inflow’ model of Chiappini, Matteucci & Gratton (1997). Thick curves: sketch of suggested separate AMRs for the Thick and Thin disks. Adapted from Chiappini et al. (1997).

This gives further support to a short time-scale for Thick-disk star formation, confirmed by HIPPARCOS parallaxes (Ng & Bertelli 1998), which show a hiatus between that and the first stars of the Thin disk, also visible in the data from Edvardsson et al. (1993) shown in Fig 3. The need for two time-scales and a hiatus in star formation, as postulated by Chiappini et al., is clearly evident, but their model does not go far enough, and it seems more realistic to draw in two entirely separate age-metallicity relations as shown schematically by the thick curves.

We also face the complication that the iron-to- α -element conversion is different in the two disks. Making use of the relationships embodied in Fig 2, we can convert Fig 1 into the Mg MDF shown in Fig 4.

Fig 4 shows some interesting features not previously apparent. For one thing, the upper abundance limits are identical and the means nearly so, so there is no longer any basis for assuming Hartwick-style mass outflow from the Thick disk with a resulting reduction in effective yield. It does have a low-metallicity tail, also noticed by Beers & Sommer-Larsen (1995), whereas the Thin one has a sharp lower limit suggestive of PIE; but if the previous




Figure 4. Magnesium MDFs. Histograms: data after Wyse & Gilmore (1995). Curves: dashed, simple inflow model for Thick disk; dotted, PIE plus inflow model for Thin disk; dash-dotted, inflow model without PIE; long-dash dotted, PIE without inflow.

metallicity came from the Thick disk I would rather call it Prompt Initial Impoverishment!

The curves in Fig 4 show some analytical fits to the two MDFs. The Thick disk is readily fitted by a simple decaying inflow model as in Pagel (1997) with a yield 0.15 dex below solar, a final gas fraction of 0.1 and any time scale less than that of SNIa, say around 1 Gyr.

For the Thin disk there are more choices. A pure PIE model without inflow gives a barely acceptable fit, while a pure inflow model (Clayton 1985 standard model with $k = 4$) produces too much of a low-metallicity tail. A combined PIE-inflow model fits very well, possibly representing terminal Thick-disk gas diluted by inflow of unprocessed material before the Clayton episode — thus a ‘three-inflow’ model.

3 Implications for radio-active cosmochronology

Solar-system actinide abundances have been used in the past to guess the age of the Galaxy, although there is also a clear dependence on chemical evolution models. More recently, with the discovery of r-process rich halo stars, it has become possible to use stellar thorium abundances in a manner that is less model-dependent, but the age of the halo does depend explicitly or implicitly on the Th/Eu production ratio. This has been estimated on the basis of nuclear physics, but with a fair degree of uncertainty (see Westin et al. 2000), while Truran (1999) has appealed to the proto-solar ratio, but this is clearly only a lower limit, thereby giving just a minimum possible age for the halo stars.

Table 1. K -ratios of actinide abundances at Solar System formation

	235, 238	232, 238	Eu, Th
‘Observed’ after Fowler (1987)	$0.25 \pm .04$	$1.35 \pm .10$	
Modified Fowler model	0.24	1.45	1.23
Inflow model (Clayton $k = 4$)	0.49	1.13	1.06

Now we know more about stellar ages and the gap between the two disks, it may be of interest to turn the argument round and check for consistency. We use our Galactic models to estimate the K -ratios defined by Fowler (1987):

$$K_{ij} \equiv \frac{N_i}{N_j}(\text{protosolar}) / \frac{p_i}{p_j} \quad (1)$$

and use the resulting $K_{\text{Eu Th}}$ to predict the production ratio from the known proto-solar abundance ratio.

We recall Fowler’s own model, a closed PIE model from which he deduced an age for the disk of just 10 Gyr. If we think only of the Thin disk, then this was remarkably prescient, but now we also have to think of the Thick disk or bulge where any PIE would probably have originated and which is 2 or 3 Gyr older. We assume the Thick disk to have formed instantaneously 12 Gyr ago and the Thin disk to have started forming after an interval $\epsilon = 3$ Gyr. The Solar System was formed after a further interval $\Delta = 4.4$ Gyr. Fowler’s equations are then modified to:

$$N_i(\text{radioactive}) \propto p_i \left[\frac{S}{1-S} e^{-\lambda_i(\Delta+\epsilon)} + \frac{1-e^{-\lambda_i\Delta}}{\lambda_i\Delta} \right] \quad (2)$$

$$N_k(\text{stable}) \propto p_k / (1-S) \quad (3)$$

where in our case the PIE contribution $S = 0.4$ rather than 0.17. Corresponding formulae for the inflow model have been given by Clayton (1985). Resulting K -ratios in the Solar System are given in Table 1, where it is seen that the modified Fowler model still gives excellent agreement with ‘observed’ K -ratios, whereas the inflow model gives numbers that are barely compatible if at all and presumably represents a generous upper limit to the modifications from inflow.

Implications for halo ages are given in Table 2. Fowler’s model implies a large correction from protosolar to production ratio and hence a large (and perhaps unacceptable) figure for the halo age, while the inflow model gives only a minimal correction and a more acceptable age range. It is intriguing that the range of production ratios permitted by Galactic chemical evolution models matches so closely the range deduced from nuclear physics.

Table 2. Th/Eu ratios and halo age

	Th/Eu ratio	halo age (Gyr)
Halo stars	0.22 to 0.25	
Proto-solar	0.46	
Nuclear theory prediction	0.48 to 0.55	13.2 to 18.6
Modified Fowler model prediction	0.57	16.7 to 19.3
Inflow model prediction	0.49	13.7 to 16.3

References

1. Beers, T.M. & Sommer-Larsen, J. 1995, ApJS, 96, 175
2. Chiappini, C., Matteucci, F. & Gratton, R. 1997, ApJ, 477, 765
3. Clayton, D.D. 1985,, in W.D. Arnett & J. Truran (eds), *Nucleosynthesis: Challenges and New Developments*, University of Chicago Press, p. 65
4. Edvardsson, B. et al. 1993, A & A, 275, 101
5. Fowler, W.A. 1987, QJRAS, 28, 87
6. Fuhrmann, K. 1998, A & A, 338, 161
7. Gilmore, G. & Wyse, R.F.G. 1986, Nature, 322, 806
8. Gratton, R. et al. 2000, A & A, in press, astro-ph/0004157
9. Hartwick, F.D.A. 1976, ApJ, 209, 418
10. Köppen & Arimoto, N. 1990, A & A, 240, 22
11. Larson, R.B. 1972, Nature Phys. Sci., 236, 7
12. Matteucci, F. & François, 1989, MNRAS, 239, 885
13. Ng, Y.K. & Bertelli, G. 1998, A & A, 329, 943
14. Pagel, B.E.J. 1989, Rev. Mex. Astr. Astrofis., 18, 153
15. Pagel, B.E.J. 1997, Nucleosynthesis and Chemical Evolution of Galaxies, Cambridge University Press, p. 243
16. Pagel, B.E.J. & Tautvaišienė, G. 1995, MNRAS, 276, 505
17. Pardi, M.C., Ferrini, F. & Matteucci, F. 1995, ApJ, 444, 207
18. Samland, M., Hensler, G. & Theis, Ch. 1997, ApJ, 476, 544
19. Strobel, A. 1991, A & A, 247, 35
20. Truran, J.W. 1999, in R. Diehl & D. Hartmann (eds), *Astronomy with Radioactivities*, Ringberg Workshop, MPE Rep. 274, p. 261
21. Truran, J.W. & Cameron, A.G.W. 1971, Ap. Sp. Sci., 14, 179
22. Westin, J., Sneden, C., Gustafsson, B. & Cowan, J.J. 2000, ApJ, 530, 783
23. Wyse, R.F.G. & Gilmore, G. 1992, AJ, 104, 144
24. Wyse, R.F.G. & Gilmore, G. 1995, AJ, 110, 2771